

Concerned about opioids after
C-section surgery?

THINK ABOUT NON-OPIOID PAIN RELIEF

Ask your
doctor about
EXPAREL for
quicker recovery
after surgery

Please see Indication and Important Safety Information on
reverse and refer to accompanying full Prescribing Information.

For more information, please visit www.EXPAREL.com/patient
or call 1-855-793-9727.

NON-OPIOID
EXPAREL[®]
(bupivacaine liposome injectable suspension)

 >10 MILLION PATIENTS HAVE RECEIVED EXPAREL

EXPAREL is a long-lasting numbing medicine that can improve your recovery after surgery

- EXPAREL is a non-addictive, non-opioid medicine
- EXPAREL can be used for women who plan to breastfeed
- EXPAREL works directly where you had your surgery, with pain relief lasting up to 3 days*
- EXPAREL has been proven to reduce length of stay in the hospital, allowing you to recover in the comfort of your own home
 - In C-section studies, women who received EXPAREL left the hospital 1 day sooner
- EXPAREL may reduce your need for pain medicine such as opioids[†]

EXPAREL is given during surgery.
Be sure to talk to your doctor about
EXPAREL *before* your surgery.



Scan here to
learn more

*Based on clinical studies of patients who had C-sections and received EXPAREL compared with patients who did not receive EXPAREL.

[†]The clinical benefit of the decrease in opioid consumption was not demonstrated in the pivotal trials.

Indication

EXPAREL is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Important Safety Information

EXPAREL should not be used in obstetrical paracervical block anesthesia.

In studies in adults where EXPAREL was injected into a wound, the most common side effects were nausea, constipation, and vomiting.

In studies in adults where EXPAREL was injected near a nerve, the most common side effects were nausea, fever, and constipation.

In the study where EXPAREL was given to children, the most common side effects were nausea, vomiting, constipation, low blood pressure, low number of red blood cells, muscle twitching, blurred vision, itching, and rapid heartbeat.

EXPAREL can cause a temporary loss of feeling and/or loss of muscle movement. How much and how long the loss of feeling and/or muscle movement depends on where and how much of EXPAREL was injected and may last for up to 5 days.

EXPAREL is not recommended to be used in patients younger than 6 years old for injection into the wound, for patients younger than 18 years old for injection near a nerve, and/or in pregnant women.

Tell your health care provider if you or your child has liver disease, since this may affect how the active ingredient (bupivacaine) in EXPAREL is eliminated from the body.

EXPAREL should not be injected into the spine, joints, or veins.

The active ingredient in EXPAREL can affect the nervous system and the cardiovascular system; may cause an allergic reaction; may cause damage if injected into the joints; and can cause a rare blood disorder.

Please refer to accompanying full Prescribing Information.

For more information, please visit www.EXPAREL.com/patient or call 1-855-793-9727.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.



©2022 Pacira Pharmaceuticals, Inc., a wholly owned subsidiary of Pacira BioSciences, Inc. All rights reserved. PP-EX-US-7676 06/22

NON-OPIOID
EXPAREL[®]
(bupivacaine liposome injectable suspension)