

Administration Case Report With EXPAREL

This case report represents the individual experience of Dr Charles Miller and is intended to demonstrate his methodology for using EXPAREL in a specific soft tissue surgery.

Pacira Pharmaceuticals, Inc. recognizes that there are alternative methodologies for administering local anesthetics, as well as individual patient considerations, when selecting the dose for a specific procedure.

EXPAREL is indicated for single-dose infiltration in adults to produce postsurgical local analgesia and as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.


CASE INFORMATION

Physician Name	Charles Miller, MD
Affiliation	Advocate Lutheran General Hospital; Park Ridge, IL
Surgical Case Performed	Robotic-assisted, laparoscopic hysterectomy, with bilateral salpingectomy
Inpatient or Outpatient Procedure	Outpatient

PATIENT CHARACTERISTICS

Gender	Female
Age	36 years of age
Patient History and Characteristics	History of severe stage IV endometriosis and dysmenorrhea

PROCEDURAL DETAILS

Incision Size	One 10-mm incision at the umbilicus; two 8-mm lateral incisions
Preoperative Analgesics Used	Acetaminophen 975 mg, pregabalin 150 mg, and celecoxib 200 mg administered 2 hours before surgery with a sip of water
Intraoperative Analgesics Used	0.5% bupivacaine HCl (<10 mL)
Dose of EXPAREL and Total Volume Used	 EXPAREL (266 mg) + Normal Saline = Total 40 mL

The recommended dose of EXPAREL is based on the size of the surgical site, the volume required to cover the area, and individual patient factors that may impact the safety of an amide local anesthetic. The maximum dose of EXPAREL should not exceed 266 mg.

EXPAREL can be administered unexpanded (20 mL) or expanded to increase volume up to a total of 300 mL (final concentration of 0.89 mg/mL [ie, 1:14 dilution by volume]) with normal (0.9%) saline or lactated Ringer's solution.

Bupivacaine HCl may be administered immediately before EXPAREL or admixed in the same syringe, as long as the ratio of the milligram dose of bupivacaine HCl to EXPAREL does not exceed 1:2. Admixing may impact the pharmacokinetic and/or physiochemical properties of EXPAREL, and this effect is concentration dependent. The toxic effects of these drugs are additive and their administration should be used with caution, including monitoring for neurological and cardiovascular effects related to local anesthetic systemic toxicity. Other than with bupivacaine, EXPAREL should not be admixed with other drugs prior to administration.

Please see Important Safety Information on reverse and refer to the accompanying full Prescribing Information before using EXPAREL for complete Dosage and Administration information.

INFILTRATION NOTES

- Using a 10-mL, 19-gauge needle:
 - 10 mL of EXPAREL® (bupivacaine liposome injectable suspension) solution was infiltrated into each lateral port down to the level of the fascia in a circumferential fashion for a total volume of 20 mL (see Figure 1)
 - 20 mL of EXPAREL solution was infiltrated around the umbilicus down to the level of the fascia in a circumferential fashion (see Figure 2)
- One-layer closures were performed on each port following the infiltration



Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia. Adverse reactions reported with an incidence greater than or equal to 10% following EXPAREL administration via infiltration were nausea, constipation, and vomiting; adverse reactions reported with an incidence greater than or equal to 10% following EXPAREL administration via interscalene brachial plexus nerve block were nausea, pyrexia, and constipation. If EXPAREL and other non-bupivacaine local anesthetics, including lidocaine, are administered at the same site, there may be an immediate release of bupivacaine from EXPAREL. Therefore, EXPAREL may be administered to the same site 20 minutes after injecting lidocaine. EXPAREL is not recommended to be used in the following patient population: patients <18 years old and/or pregnant patients. Because amide-type local anesthetics, such as bupivacaine, are metabolized by the liver, EXPAREL should be used cautiously in patients with hepatic disease.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL. EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use. The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression. **Cardiovascular System Reactions:** Toxic blood concentrations depress cardiac conductivity and excitability which may lead to dysrhythmias, sometimes leading to death. **Allergic Reactions:** Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients. **Chondrolysis:** There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Disclosure: Dr Miller has no relationship with Pacira to disclose.